Precautions After Belody Lifting Treatments

- Swelling, a sensation of heat, bruising, and other reactions may occur depending on the area treated or the individual's body constitution.
- Continuing with ice pack massage for about 5 days after the procedure can help reduce swelling, and it is important to take prescribed antibiotics to prevent infection. (After 5 days, please switch to warm packs for the best recovery.)
- A sensation of tightness may last for about one week on average.
- You can shower or wash the treated area the day after surgery, but you must apply a waterproof bandage over the incision site before doing so. (Please remove the waterproof bandage immediately after showering)
- Avoid alcohol and smoking until swelling and bruising have subsided.
- For about a week after the procedure, you should avoid saunas, steam rooms, and swimming.
- On rare occasions, complications such as infections, bleeding, or foreign body reactions may occur at the treated site.
- Stitch removal will be performed 7 days after facial procedures and 14 days after body procedures.

※ Individual experiences may vary.

This Page

WhatsApp

SEOJIN
PLASTIC SURGERY

Email: contact@seojinps.co
WhatsApp: +82 10 3555 5167
Address: 8F, EGI Building, 843 Nonhyeon-ro (Sinsa-dong), Gangnam-gu, Seoul

Free Online
Consultation

Company name: Seojin Plastic Surgery Clinic | Representative: Hyungmin Lee | Business registration number: 588-31-01105

Copyright © 2026 Seojin PS